Sadbovezemiaky.sk

Variety sheet

Paradiso



A reliable variety with impressive yields, a high number of tubers and attractive potatoes. A strong plant that also performs well in warmer and drier areas.

Parentage **Rodeo × Valor**

Note: The breeder lists Paradiso as early maincrop; based on our experience we classify it as an early variety.

Cooking type B	Maturity	Use	Tubers	Skin and flesh	Storability
Fairly firm, all-purpose	**Early**	**Table use**	**Large, long-oval**	**Yellow / light yellow**	**Medium**

Tubers and quality

Tuber shape	**Long-oval**
Tuber size	**Large**
Number of tubers	**Medium**
Shape uniformity	**Fairly uniform**
Skin finish	**Good**
Dormancy	**Medium**
Dry matter	**18.2 %**
Underwater weight	**329 g** (UWW)
Discolouration after cooking	**Slight**

Resistance

1 lowest, 9 highest

Late blight, foliage	6
Late blight, tubers	6
Common scab	6
Powdery scab	8
Virus Y (PVY)	6
Tuber necrosis (Yntn)	8
Spraing (TRV)	4
Potato cyst nematode Ro1, Ro4	**Resistant**
Wart disease pathotype 1	**Resistant**

Why Paradiso

- Impressive and stable yield
- High number of tubers
- Stable yield even in drier areas
- Attractive, uniform tubers

In the field

Initial foliage development	**Fairly fast**
Metribuzin (Sencor)	**Not sensitive pre-, fairly post-emergence**
Mechanical damage	**Fairly sensitive**
Internal bruising	**Slightly sensitive**
Growth cracks	**Fairly sensitive**
Secondary growth	**Slightly sensitive**

Data are based on the breeder's information and our own experience. They are indicative; results always depend on soil, weather and crop management.

Agriana s.r.o., Mila Urbana 18, 957 01 Bánovce nad Bebravou
objednavky@sadbovezemiaky.sk, +421 38 760 2398, +421 911 920 398

Cultivation

From planting to storage

1 Seed preparation and planting

- Better not to remove sprouts; it reduces yield and uniformity.
- Plant into warm soil, ideally at least 8 °C.

2 During the growing season

- Split the nitrogen; give the second dose shortly after tuber set.
- Metribuzin (Sencor): not sensitive pre-emergence, fairly sensitive post-emergence.
- Preventive late blight protection is recommended.

3 Haulm killing and harvest

- Harvest no sooner than 2 weeks after haulm killing; tubers are more prone to damage.

4 Storage medium-term

- Suitable for medium-term storage.
- Recommended storage temperature 4 °C.



Seed potatoes from us

Availability	on order
Orders	by e-mail or phone

Order Paradiso
Scan the code. We ship to Slovakia and the Czech Republic.

Planting density

Seed size	Plants per ha	In-row spacing
28 – 35 mm	58,000	23 cm
35 – 55 mm	38,000	35 cm

Rows 75 cm. Always check the number of tubers per 1 kg of seed.

Fertilisation

N	P	K
−10 %	standard rate	standard rate
compared to the standard rate, split		

Always adjust rates to soil analysis.